

2016

## Session D, 2016 First Place: Mighty Moss: The water-holding capacity, heat retention, and debris content of local moss genera in relation to its use as a traditional diaper material

Jodie Schoelkopf, Emma Livingston, Mia Eddy, Megan Ferreira

Follow this and additional works at: <https://digitalcommons.esf.edu/clbs>



Part of the [Plant Sciences Commons](#)

---

### Recommended Citation

Jodie Schoelkopf, Emma Livingston, Mia Eddy, Megan Ferreira, "Session D, 2016 First Place: Mighty Moss: The water-holding capacity, heat retention, and debris content of local moss genera in relation to its use as a traditional diaper material" (2016). *Cranberry Lake Biological Station*. 16.  
<https://digitalcommons.esf.edu/clbs/16>

This Presentation is brought to you for free and open access by the Environmental and Forest Biology at Digital Commons @ ESF. It has been accepted for inclusion in Cranberry Lake Biological Station by an authorized administrator of Digital Commons @ ESF. For more information, please contact [digitalcommons@esf.edu](mailto:digitalcommons@esf.edu), [cjkoons@esf.edu](mailto:cjkoons@esf.edu).

# Mighty Moss:

## The water-holding capacity, heat retention, and debris content of local moss genera in relation to its use as a traditional diaper material

---

Sphagnum Sisters

Jodie Schoelkopf, Megan Ferreira, Emma Livingston,  
& Mia Eddy



# Rationale & Importance of the Study

- Vast amounts of indigenous knowledge, but limited amount of literature that connects traditional plant usage with Western scientific methodology (Harris 2008)
- Mosses have been used in different cultures worldwide (Harris 2008 and Glime 2013)
- What moss genus would have been used locally as diaper material? Why?



Photo by Bob Klips

# Chosen Moss Genera (Experimental Units)

Carpet Moss  
(*Dicranum*)



- Insect repellent
- Resistant to mold
- Diaper lining (Johnson 1995)

Haircap Moss  
(*Polytrichum*)



- Cloaks (Beever 1995)

Pincushion Moss  
(*Leucobryum*)



- Cushion and mattress stuffing (Glime 2013)

Peat Moss  
(*Sphagnum*)




- Cradles, Diapers
- Insulation
- Pillow and mattress stuffing (Harris 2008)

Processed Cotton  
(*Gossypium*)



- Modern day diapers (Mesek 1981)

# Collection of Moss



Cranberry Lake

| | = 5000 ft

# Data

- Categorical
- Quantitative

# Statistical Tests

- Descriptive statistics
- ANOVA

# Sample Units

- Experiment 1: 10g fresh samples of each experimental unit
- Experiment 2: 10g samples of each experimental unit, saturated in water
- Experiment 3: Mesh bags of each experimental unit, density of 1g/mL

# Hypothesis 1: Contained Debris

$H_{01}$ : There will be no difference between the amount of debris (g) filtered from each moss

$H_{a1}$ : *Sphagnum* will contain the least debris (g) relative to other moss genera

# Methods:

- 10g samples agitated in water and debris allowed to fall to bottom
- Debris was filtered through filter paper
- The debris covered filter paper was dried in a drying oven
- 10 trials



Carpet


Pincushion

Sphagnum

Haircap



# Results: Contained Debris



F-value = 7.93

P-value = 0.0003

SE = 0.3676

DF = 3

Figure 1. Debris entrapped within 10g of each variable. Variable grouped using Tukey's test.

# Hypothesis 2: Water Holding Capacity

H<sub>02</sub>: There will be no difference between the water holding capacity (mL) of mosses


H<sub>a2</sub>: *Sphagnum* will have the highest water holding capacity (mL) relative to other moss genera

# Methods: Water Capacity

- Substrates soaked for ten minutes to allow maximum absorption
- Dehydrated until brittle
- Dry weight (g) subtracted from wet weight to calculate the amount of water (mL) retained in each variable
- 10 trials



# Results: Water Capacity



$F = 75.83$   
 $P\text{-value} = <0.001$   
 $SE = 0.1619$   
 $DF = 4$

Figure 2. Amount of water (mL) in 10g of saturated substrate. Variables grouped using Tukey's test.

## Hypothesis 3: Heat Retention

$H_{03}$ : There will be no difference in the rate of heat loss ( $^{\circ}\text{C}/\text{min}$ ) between all variables

$H_{a3}$ : *Sphagnum* will have the slowest rate of heat loss ( $^{\circ}\text{C}/\text{min}$ ) relative to the other variables

# Methods:

Each material placed in a mesh bag at a density of 1g/mL

- The mesh bags were heated by being soaked in 37.7 °C water for five minutes
- Removed from heat
- Temperature change tracked every five minutes over a span of 60 minutes.



# Results: Heat Retention




Figure 3. Heat loss over time (°C/min) in each variable over 60 minutes.




Figure 4. Mean heat loss over time (°C/min) in each variable. Variables grouped with Tukey's test.

F-value = 2.32  
P-value = 0.091  
SE = 0.01665  
DF = 4

# Discussion: Reasoning Behind Results

- *Sphagnum* made up of a single layer of cells
  - Photosynthetic cells and dead hyaline cells (Glime 2013)
  - Whorled structure of *sphagnum* and the absorbent hyaline cells allow for high water capacity
  - An osmophillic layer with pores for absorption (Glime 2013)
- *Polytrichum* has both water and sugar conducting cells
  - can obtain nutrients from the soil rather than the environment (Glime 2013).
  - also have waxy cuticle that repels rather than absorbs water (Proctor, 1979).



# Discussion: Why does it matter?

- Supports Traditional Ecological Knowledge (TEK)
- Indigenous peoples in the region would have most likely chosen *Sphagnum* for use in diapers
- Comparable to today's cotton diapers
  - ... in fact it's better!
  - Biodegradable and unprocessed



Photo by Megan Ferreira 2016

# Follow Up Studies

- Abundance of moss genera in different regions to infer which may have been used most by the people of that area
- Antimicrobial properties (Singh et. al., 2007 and Hotson, 1921)
  - Evolutionary relationship between moss usage in nest building and fecal sacs in birds
- Heat retention in samples of non-saturated moss genera
  - Water may have acted as an insulator



Photo by Jay Greenberg



Photo by Megan Ferreira 2016

# Conclusion

- *Sphagnum* contained the least amount of debris
  - Water retention is greatest in *Sphagnum*, on par with processed cotton
  - Heat retention is greatest in *Sphagnum* compared to cotton and other moss genera tested, however is not statistically significant
  - *Polytrichum* did not retain water and heat well, and contained the most debris
- This data supports the use of *Sphagnum* by traditional cultures in its cleanliness and absorbent properties, and suggests it has the greatest potential as a traditional diaper material.

# Acknowledgments

- Dr. Robin W. Kimmerer
- Dr. Kim Shultz
- Dr. Melissa Fierke
- Mr. Giuseppe Tumminello
- Work studs: Joey, Michelle, & Kristen
- The moss

Thank you!



Photo by Bob Klips

# References

- Beever, J. E. & Gresson, J. E (1995). *Polytrichum commune* Hedw. and *Polytrichadelphus magellanicus* (Hedw.) Mitt. used as decorative material on New Zealand Maori cloaks. *The Bryologist*, 18(4), 819-823. doi: 10.1179/jbr.1995.18.4.819.
- Glime, J. M. (2013). pH Lowering Ability of Sphagnum. *Bryophyte Ecology* (Vol. 5). Houghton, MI.
- Harris, E. S. (2008). Ethnobryology: Traditional uses and folk classification of bryophytes. *The Bryologist*, 111(2), 169-217.
- Hotson, J. W. (1921). Sphagnum Used as a Surgical Dressing in Germany during the World War. *The Bryologist*. 24(5), 74. doi:10.2307/3238003
- Johnson Gottesfeld, L. M. and Vitt, D. H. 1996. The selection of Sphagnum for diapers by indigenous North Americans. *Evansia* 13: 103-108.
- Mesek, Frederick. 1981. "Disposable Diaper" Patent 4285342
- Proctor, M. C. (1979). Surface wax on the leaves of some mosses. *Journal of Bryology*, 10(4), 531-538. doi:10.1179/jbr.1979.10.4.531
- Singh, M., Rawat, A., & Govindarajan, R. (2007). Antimicrobial activity of some Indian mosses. *Fitoterapia*, 78(2), 156-158. doi:10.1016/j.fitote.2006.10.008

**THIS IS YOUR BABY ON PAMPERS.**





# THIS IS YOUR BABY ON SPHAGNUM.



Photo by Doug Elliott

# Questions?



Photo by Doug Elliott